



Emulsification technique affects oxidative stability of fish oil-in-water emulsions

Horn, Anna Frisenfeldt; Nielsen, Nina Skall; Jensen, Louise Helene Søggaard; Horseywell, Andy; Jacobsen, Charlotte

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Horn, A. F., Nielsen, N. S., Jensen, L. H. S., Horseywell, A., & Jacobsen, C. (2012). *Emulsification technique affects oxidative stability of fish oil-in-water emulsions*. Abstract from Science and Technology of Food Emulsions, London, United Kingdom.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Conference abstract (Poster presentation)

Science and Technology of Food Emulsions, June 21-22, 2012, London, UK

Emulsification technique affects oxidative stability of fish oil-in-water emulsions

Anna Frisenfeldt Horn¹, Nina Skall Nielsen¹, Louise Søgaaard Jensen^{2,3}, Andy Horseywell^{2,3} & Charlotte Jacobsen¹

¹National Food Institute, Technical University of Denmark

²Center for Electron Nanoscopy, Technical University of Denmark

³Department of Mechanical Engineering, Technical University of Denmark

In oil-in-water emulsions, lipid oxidation is expected to be initiated at the oil-water interface. The properties of the emulsifier used, and the structure at the interface is therefore expected to be of great importance for lipid oxidation in emulsions. Previous studies have shown that e.g. homogenization pressure can affect how proteins locate themselves at the interface of an emulsion. The hypothesis is therefore that emulsions produced with different emulsification equipments differ in their oxidative stability due to differences in the behaviour of the proteins at the interface.

The aim of this study was to compare lipid oxidation in 10% fish oil-in-water emulsions prepared by two different kinds of high pressure homogenizers i.e. a microfluidizer and a two valve high pressure homogenizer. Emulsions were made with equal droplet sizes, and with either 1% sodium caseinate or 1% whey protein isolate. Emulsions were characterised and investigated by microscopy. Lipid oxidation was assessed by PV and the formation of secondary volatile oxidation products. Results showed that the different emulsification techniques had an influence on lipid oxidation and that the effect of the emulsification technique depended on the type of protein used as an emulsifier.